




## Kentville Research & Development Centre (KRDC) – Nova Scotia wine grape bud hardiness 2025/2026 Report no. 4: February 18 & 19, 2026

Prepared by Jeff Franklin ([jeff.franklin@agr.gc.ca](mailto:jeff.franklin@agr.gc.ca)) and Dr. Harrison Wright ([harrison.wright@agr.gc.ca](mailto:harrison.wright@agr.gc.ca)), Plant Physiology Program, KRDC, Agriculture and Agri-Food Canada (AAFC) / Government of Canada; 32 Main St, Kentville, Nova Scotia, B4N 1J5.



**Figure 1.** Plot showing the LTE50 values (coloured lines) for four wine grape varieties taken from Nova Scotia vineyards, as well as recent and historical temperature trends. Current observed minimum temperatures (black line) as well as the 100-year minimum temperatures (grey line) were recorded at the Environment and Climate Change Canada (ECCC) weather station located at the Kentville Research and Development Centre.



### Current report

In this round of the survey, we find that most of the varieties are staying close to their deep winter minimum LTE50 values. The exception to this is New York Muscat, which has deacclimated slightly (1.3 °C). The reason for this is not clear as we would expect all varieties to stay at their deep winter values for another week or so given the February temperatures that we have experienced. So far in February, the mean daily temperature has been -4.3 °C, which is equal to the 25-year average. The forecast for the next 14 days calls for another week of below average temperatures followed by a period of above average temperatures. At this time, the Environment and Climate Change Canada 30-day forecast suggests that the southern half of Nova Scotia will experience above average temperatures for the month of March. Although it is too early to make solid predictions, it appears that spring may not be too far away.

**Table 1.** LTE10, LTE50 and LTE90 average values (°C) for core wine grape cultivars for the current and previous reporting periods

Core cultivars and sites	Nov. 4 - 5			Jan. 6 - 7			Feb. 4 - 5			Feb. 18 -19			LTE10	LTE50	LTE90
	LTE10	LTE50	LTE90	LTE10	LTE50	LTE90	LTE10	LTE50	LTE90	LTE10	LTE50	LTE90			
Chardonnay (5 sites)	-14.0	-16.1	-18.9	-22.6	-24.4	-25.8	-23.2	-24.8	-26.0	-22.9	-24.3	-25.6			
L'Acadie (5 sites)	-12.2	-15.2	-20.2	-24.2	-26.2	-28.0	-24.2	-27.1	-29.1	-26.2	-28.4	-30.4			
Marquette (5 sites)	-16.0	-20.2	-23.9	-25.8	-27.7	-29.1	-26.7	-28.3	-30.6	-27.2	-28.4	-30.0			
New York Muscat (5 sites)	-15.4	-18.6	-21.6	-23.6	-26.3	-27.8	-24.9	-26.6	-29.0	-23.7	-25.3	-27.2			
NYUS.2.1 LTE50 prediction*															
Chardonnay (Kentville)		-18.6			-24.6			-23.1			-23.3				
L'Acadie (Kentville)		-19.0			-29.5			-28.2			-28.0				
Marquette (Kentville)		-19.0			-27.5			-26.5			-26.1				

\* Wang et al., 2024. *Horticulture Research*, 11, 2: uhad286. Follow predicted bud hardiness values in real time at a weather site near you in the US or Canada via the following website: [https://cornell-tree-fruit-physiology.shinyapps.io/North\\_America\\_Grape\\_Freezing\\_Tolerance/](https://cornell-tree-fruit-physiology.shinyapps.io/North_America_Grape_Freezing_Tolerance/).



### Research report description

The Nova Scotia wine grape bud hardiness survey generates reports detailing the low temperature exotherm (LTE) values over the dormant period (roughly from November to April). The LTE is the temperature (°C) at which a bud freezes and is killed: LTE10, LTE50 and LTE90 values denote the temperatures at which 10%, 50% and 90% of the viable buds freeze. The LTE values for a given variety and site are generated using eight canes obtained from eight vines; the compound buds from nodes 3 through 7 from each cane are measured via differential thermal analysis (DTA). It is important to note that the LTE value denotes a bud's susceptibility to acute, cold temperature damage; it does *not* necessarily reflect the bud's susceptibility to dehydration, poor vine health and other more chronic forms of stress that can result in bud mortality at temperatures above the LTE values.

Each report includes: (1) a plot showing the median LTE50 values for a group of hybrid and vinifera wine grape cultivars averaged over several sites located in Kings county as well as recent and historical minimum temperature trends (Figure 1); (2) comments on the current reporting period; (3) a table of LTE10, LTE50 and LTE90 values for the same cultivars shown in Figure (Table 1); (4) A computer-model generated approximation of the LTE50 value based on temperatures obtained from the Kentville Environment and Climate Change Canada (ECCC) weather station and the NYUS.2 machine learning model. This report is produced by the KRDC Plant Physiology Program and is supported by Grape and Wine Cluster Activity #18: Growing More Resilient and Hardy Wine Grapes in the Face of Climate Change in an Eastern Canadian Environment. If you have any questions or comments, please feel free to reach out to the KRDC Plant Physiology Program using the contact information listed above.

© His Majesty the King in Right of Canada, represented by the Minister of Agriculture and Agri-Food (2026).